

+4 or trisomy 4

Clinics and Pathology

- Disease** Acute non lymphocytic leukaemia (ANLL)
- Epidemiology** +4 as the sole anomaly is a rare chromosomal abnormality associated with a specific subtype of primary ANLL and secondary (treatment related) ANLL with myelomonocytic morphology; it has been found with the same frequency in the M1-M2 and M4 French-American-British (FAB) phenotypes
- association of +4 with double minute chromosomes has been described in ten cases; five with AML-M2, two with AML-M4, one with refractory anemia with excess of blasts in transformation (RAEB-T), one with chronic myelomonocytic leukemia (CMMoL) and one with unclassified preleukemia
- the coincidence of +4 with t(8;21) or its variant t(6;21;8), observed in at least two cases of ANLL (M1 and M2), is therefore recurrent
- Prognosis** apparently +4 has no prognostic significance in ANLL; with the exception of the cases bearing [c-Kit](#) mutations who are associated with a rapid disease progression
- Disease** Acute lymphocytic leukaemia (ALL)
- Epidemiology** +4 has been described in two cases of T-cell acute lymphoblastic leukemia as the sole chromosomal anomaly
- combined trisomies of chromosomes 4 and 10 are found in children with B-progenitor cell acute lymphoblastic leukemia with a favourable prognostic association
- Prognosis** patients with chromosomes 4 and 10 trisomies have an extremely favourable 4-year event-free survival (EFS) after antimetabolite-based chemotherapy
- Disease** Thecoma of the ovary
- Epidemiology** +4 associated with +12 has been described in two cases of fibrothecoma and in one case of thecoma; it has been suggested that acquisition of trisomy 4 may constitute a second cytogenetic step in tumor progression of ovarian thecoma/fibrothecoma.
- Prognosis** no prognostic significance

Disease +4 has been found as the sole aberration only in one case of [uterine leiomyoma](#)

Genetics

trisomy 4 was proven to lead to duplication of Asp816Tyr mutation of c-Kit gene (that maps to 4q12) in a case of M2-ANLL with mast cell involvement

External links

Other database [+4 or trisomy 4](#) [Mitelman database \(CGAP - NCBI\)](#)

Bibliography

Trisomy 4 identifies a subset of acute nonlymphocytic leukemias.

Mecucci C, Van Orshoven A, Tricot G, Michaux JL, Delannoy A, Van Den Berghe H
Blood 1986; 67: 1328-1332.

Trisomy of leukemic cell chromosomes 4 and 10 identifies children with B-Progenitor cell acute lymphoblastic leukemia with a very low risk of treatment failure: a pediatric oncology group study.

Harris MB, Shuster JJ, Carrol A, Look AT, Borowitz MJ, Crist WM, Nitschke R, Pullen J, Steuber CP, Land VJ.
Blood 1992; 79: 3316-3324.

Trisomy 12 and 4 in a thecoma of the ovary.

Mrozek K, Limon J, Debniak J, Emerich J.
Gynecol Oncol. 1992; 45: 66-68.

Trisomy 4 may occur in a broad range of hematologic malignancies.

Kwong YL, Ha SY, Liu HW, Chan LC.
Cancer Genet Cytogenet. 1993; 69: 139-140.

Trisomy 4 in stem cell like leukemic cells of a patient with AML.

Feuring-Buske M, Haase D, Konemann S, Troff C, Grove D, Hiddemann W, Wormann B.
Leukemia 1995; 9: 1318-1320.

MYC amplification in two further cases of acute myeloid leukemia with trisomy 4 and double minute chromosomes.

O'Malley F, Rayeroux K, Cole-Sinclair M, Tong M, Campbell LJ.
Cancer Genet Cytogenet 1999; 109: 123-125.

Trisomy 4 leading to duplication of a mutated KIT allele in acute myeloid

leukemia with mast cell involvement.

Beghini A, Ripamonti CB, Castorina P, Pezzetti L, Doneda L, Cairoli R, Morra E, Larizza L.

Cancer Genet Cytogenet 2000; 119: 26-31.

Contributor(s)

Written 07-
2000 Alessandro Beghini

Citation

This paper should be referenced as such :

Beghini A . +4 or trisomy 4. Atlas Genet Cytogenet Oncol Haematol. July 2000 .

URL : <http://AtlasGeneticsOncology.org/Anomalies/tri4ID1011.html>

© *Atlas of Genetics and Cytogenetics in Oncology and Haematology*
