

FCGR2B

Identity

Other names

CD32

FCG2

FCGR2

IGFR2

Hugo

FCGR2B

Location

1q22

location, by FISH/genome sequencing; FCGR2B is contained within a locus that spans approximately 200kb and that encodes 5 low affinity IgG Fc receptor genes. The published gene orientation is as follows; Cen 1q -- 3' FCGR2A 5' -- 5'- FCGR3A - 3' -- 3' FCGR2B 5' -- 5' - FCGR3B - 3' -- 5'- FCGR2C- 3'.

DNA/RNA

Description The FCGR2B gene spans a 15kb genomic region; Gene orientation : centromere - 3' FCGR2B 5' - telomere; 8 exons (denoted S1, S2, EC1, EC2, TM, IC1, IC2, IC3). S1 and S2 encode the signal peptide domain, EC1 and EC2 encode the extracellular domain, TM encodes the transmembrane domain and IC1, IC2, IC3 encode the intracytoplasmic region.

Transcription 1.7kb mRNA. Two alternately spliced mRNA exist in humans; FcgRIIb1 and FcgRIIb2 (a 19 aa insertion encoded by IC1 is present in b1 but not in b2).

Protein

Note called Fc gamma RIIB

Description 40kD transmembrane glycoprotein that possesses an intracytoplasmic Immunoreceptor Tyrosine-based Inhibition Motif (ITIM) (encoded by IC3).

Expression B, myeloid and mast cells.

Localisation Transmembrane.

Function Low affinity IgG Fc receptor. Founding member ITIM family of immune inhibitory receptors. Fc gamma RIIB functions to down-regulate activatory responses mediated by Immunoreceptor Tyrosine-based Inhibition Motif (ITAM) bearing receptors such as the B-cell receptor on B cells. For example, on B-cells, negative regulation of B-cell receptor activatory signalling is initiated through co-recruitment of BCR and Fc

gamma RIIb to IgG/ immune complexes. This leads to tyrosyl phosphorylation of the Fc gamma RIIb ITIM motif / recruitment of the inositol phosphatase SHIP and activation of the adaptor protein Dok - these events ultimately lead to the inhibition of ITAM-dependant Ca⁺⁺ mobilisation and cellular proliferation (reduced MAPK signalling), respectively.

A proapoptotic signal through the Fc gamma RIIb transmembrane sequence has been described in conditions of homo-aggregation of Fc gamma RII on murine B cells.

Homology Contains two immunoglobulin-like C2 domains.

Implicated in

Entity	1q21-23 rearrangements in NHL (Nnon-Hodgkins Lymphoma)
Disease	NHL (< 5% of cases with 1q21-23 breaks)
Prognosis	It is now known that cytogenetically determined 1q21-23 breaks can target a diversity of 1q21-23 genes. Early data has suggested a poor prognosis for 1q21 rearrangements in diffuse large cell lymphoma .
Cytogenetics	Deregulation of the FCGR2B gene by chromosomal translocation was first demonstrated in 3 NHL patients that showed a t(1;22)(q22;q22) in association with a t(14;18)(q32;q21) . Two further cases of NHL with rearrangements affecting the FCGR2B region have since been identified. Both cases also showed t(14;18)(see below for description). This suggests a role for this deregulation in lymphoma progression.
Hybrid/Mutated Gene	Dysregulation of the FCGR2B gene has been identified as a consequence of a t(1;22)(q22;q11) (3 patients) and t(1;14)(q21;q32) (1 patient). The FCGR2B coding sequence and promoter region appear to remain intact in these translocations. The principal consequence is overexpression of the Fc gamma RIIb2 isoform of Fc gamma RIIb. Deregulation of FCGR2B most likely occurs as a consequence of Ig gene transcriptional enhancer activity. Other mechanisms may exist ; an NHL patient with dup(1)(q21q25) and rearrangement in the FCGR2B region has also been identified.
Abnormal Protein	None
Oncogenesis	Contribution of FCGR2B deregulation to lymphoma development or progression is currently not known.

External links

	Nomenclature
Hugo	FCGR2B
GDB	FCGR2B
Entrez Gene	FCGR2B 2213 Fc fragment of IgG, low affinity IIb, receptor for (CD32)

Cards

[Atlas](#) [FCGR2BID397](#)

[GeneCards](#) [FCGR2B](#)

[Ensembl](#) [FCGR2B](#)

[CancerGene](#) [FCGR2B](#)

[Genatlas](#) [FCGR2B](#)

[GeneLynx](#) [FCGR2B](#)

[eGenome](#) [FCGR2B](#)

[euGene](#) [2213](#)

Genomic and cartography

[GoldenPath](#) [FCGR2B](#) - [1q22](#) [chr1:158364611-158379618](#) + [1q23.3](#) (hg17-May_2004)

[Ensembl](#) [FCGR2B](#) - [1q23.3](#) [[CytoView](#)]

[NCBI](#) [Genes Cyto](#) [Gene Seq](#) [Map View - NCBI]

[OMIM](#) [Disease map](#) [OMIM]

[HomoloGene](#) [FCGR2B](#)

Gene and transcription

[Genbank](#) [AB060143](#) [SRS] [AB060143](#) [ENTREZ]

[Genbank](#) [AB060144](#) [SRS] [AB060144](#) [ENTREZ]

[Genbank](#) [AB060145](#) [SRS] [AB060145](#) [ENTREZ]

[Genbank](#) [AB060146](#) [SRS] [AB060146](#) [ENTREZ]

[Genbank](#) [AB060147](#) [SRS] [AB060147](#) [ENTREZ]

[RefSeq](#) [NM_001002273](#) [SRS] [NM_001002273](#) [ENTREZ]

[RefSeq](#) [NM_001002274](#) [SRS] [NM_001002274](#) [ENTREZ]

[RefSeq](#) [NM_001002275](#) [SRS] [NM_001002275](#) [ENTREZ]

[RefSeq](#) [NM_004001](#) [SRS] [NM_004001](#) [ENTREZ]

[RefSeq](#) [NT_086598](#) [SRS] [NT_086598](#) [ENTREZ]

[AceView](#) [FCGR2B](#) AceView - NCBI

[TRASER](#) [FCGR2B](#) Traser - Stanford

[Unigene](#) [Hs.492655](#) [SRS] [Hs.492655](#) [NCBI] [HS492655](#) [spliceNest]

Protein : pattern, domain, 3D structure

[SwissProt](#) [P31994](#) [SRS] [P31994](#) [EXPASY] [P31994](#) [INTERPRO]

[Prosite](#) [PS50835 IG LIKE](#) [SRS] [PS50835 IG LIKE](#) [Expasy]

[Interpro](#) [IPR007110 Iq-like](#) [SRS] [IPR007110 Iq-like](#) [EBI]

[CluSTr](#) [P31994](#)

[Pfam](#) [PF00047 ig](#) [SRS] [PF00047 ig](#) [Sanger] [pfam00047](#) [NCBI-CDD]

[Blocks](#) [P31994](#)

[PDB](#) [2FCB](#) [SRS] [2FCB](#) [PdbSum], [2FCB](#) [IMB]

Polymorphism : SNP, mutations, diseases

[OMIM](#) [604590](#) [[map](#)]
[GENECLINICS](#) [604590](#)
[SNP](#) [FCGR2B](#) [dbSNP-NCBI]
[SNP](#) [NM_001002273](#) [SNP-NCI]
[SNP](#) [NM_001002274](#) [SNP-NCI]
[SNP](#) [NM_001002275](#) [SNP-NCI]
[SNP](#) [NM_004001](#) [SNP-NCI]
[SNP](#) [FCGR2B](#) [GeneSNPs - Utah] [FCGR2B](#) [SNP - CSHL] [FCGR2B](#) [HGBase - SRS]

General knowledge

[Family Browser](#) [FCGR2B](#) [UCSC Family Browser]
[SOURCE](#) [NM_001002273](#)
[SOURCE](#) [NM_001002274](#)
[SOURCE](#) [NM_001002275](#)
[SOURCE](#) [NM_004001](#)
[SMD](#) [Hs.492655](#)
[SAGE](#) [Hs.492655](#)
[Amigo](#) [function|IgG binding](#)
[Amigo](#) [process|immune response](#)
[Amigo](#) [component|integral to membrane](#)
[Amigo](#) [component|plasma membrane](#)
[Amigo](#) [function|receptor activity](#)
[Amigo](#) [process|signal transduction](#)
[BIOCARTA](#) [B Lymphocyte Cell Surface Molecules](#)
[PubGene](#) [FCGR2B](#)

Other databases

Probes

[Probe](#) [Cancer Cytogenetics \(Bari\)](#)
[Probe](#) [FCGR2B Related clones \(RZPD - Berlin\)](#)

PubMed

[PubMed](#) [18 Pubmed reference\(s\) in LocusLink](#)

Bibliography

: Fc receptor biology
 Dakron
 Annu Rev Immunol 1997;15:203-34. (Review)
 Medline [97288736](#)

: The IgG Fc receptor, FcγRIIB, is a target for deregulation by

chromosomal translocation in malignant lymphoma

Callanan MB, Le Baccon P, Mossuz P, Duley S, Bastard C, Hamoudi R, Dyer MJ, Klobeck G, Rimokh R, Sotto JJ, Leroux D.

Proc Natl Acad Sci U S A 2000 Jan 4;97(1):309-14

Medline [20087245](#)

: Immune inhibitory receptors.

Ravetch JV, Lanier LL

Science 2000 Oct 6;290(5489): 84-9 (Review)

Medline [U](#)

: Deregulation of FCGR2B expression by 1q21 rearrangements in follicular lymphomas.

Chen W, Palanisamy N, Schmidt H, Teruya-Feldstein J, Jhanwar SC, Zelenetz AD, Houldsworth J, Chaganti RS.

Oncogene 2001 Nov 15;20(52): 7686-93

Medline [21624903](#)

[REVIEW articles](#) *automatic search in PubMed*

[Last year publications](#) *automatic search in PubMed*

[BiblioGene - INIST](#)

Contributor(s)

Written 02-2002 Mary Callanan, Dominique Leroux

Citation

This paper should be referenced as such :

Callanan M, Leroux D . FCGR2B. Atlas Genet Cytogenet Oncol Haematol. February 2002 .

URL : <http://www.infobiogen.fr/services/chromcancer/Genes/FCGR2BID397.html>

© *Atlas of Genetics and Cytogenetics in Oncology and Haematology*
