

Bone tumors: an overview

Identity

Note Primary skeletal neoplasms account for 0,2% of human tumors, whereas involvement of skeletal tissue by metastatic disease is much more common. Their soft tissue-related counterparts outnumber bone tumors by a margin of approximately 10:1. Because of their rarity, not much is known about the etiology and risk factors of bone tumors, although a difference in ethnical distribution has been observed.

Bone tumors are mostly of mesenchymal origin, though for example Ewing sarcoma is thought to have neuroectodermal precursor cells. Classification of the World Health Organization will be followed in this overview. Grading of bone tumors is roughly based on the cellularity of the lesion compared to the amount of extracellular matrix, nuclear features, the presence of mitotic figures and necrosis. Staging via the TNM system is normally not used, because metastases in lymph nodes are not frequent in these lesions. Therefore staging is based on degree of differentiation of the tumor tissue and local and distant spread of the tumor.

Genetic information and references are provided for tumors investigated in more than a single case.

Classification

Cartilage tumors

- Osteochondroma
- Chondromas
- Enchondroma
- Periosteal chondroma
- Chondroblastoma
- Chondromyxoid fibroma
- Chondrosarcoma
- Dedifferentiated
- Mesenchymal
- Clear cell
- Periosteal

Osteogenic tumors

- Osteoid osteoma
- Osteoblastoma
- Osteosarcoma
- Conventional
- Teleangiectatic
- Small cell

Low grade central
Secondary
Parosteal
Periosteal
High grade surface

Fibrogenic tumors

Desmoplastic fibroma of bone
Fibrosarcoma of bone

Fibrohistiocytic tumors

Histiocytoma of bone

Ewing sarcoma/Primitive neuroectodermal tumor

Giant cell tumors

Giant cell tumor

Notochordal tumors

Chordoma

Vascular tumors

Haemangioma and related lesions
Angiosarcoma

Myogenic, lipogenic, neural and epithelial tumors

Leiomyosarcoma of bone
Lipoma of bone
Adamantinoma and osteofibrous dysplasia

Tumors of undefined neoplastic nature

Aneurysmal bone cyst
Simple bone cyst
Fibrous dysplasia
Langerhans cell histiocytosis (LCH)

Clinics and Pathology

Cytogenetics **Cartilage tumors**

Osteochondroma

Germ line mutations in the tumor suppressor genes [Exostosin-1](#) (EXT-1) located at 8q24 and [Exostosin-2](#) (EXT-2) located at 11p11-p12 have been found in hereditary multiple osteochondromas (MO). Somatic mutations are extremely rare in these tumors. In four tumors aberrations involving the region 1p13-p22 were shown.

Chondromas

- Enchondroma
By conventional cytogenetic analysis, abnormalities involving chromosome 6 and the long arm of chromosome 12 have been detected.
- Periosteal chondroma
This subtype has not been individually investigated.

Chondroblastoma

Although a dozen of case reports have been published. no specific

alterations were found in these rare benign tumors.

Chondromyxoid fibroma

Structural rearrangements of chromosome 6 are found to be non-random, particularly involving the long arm (q13 and q25) and p25 on the short arm.

Chondrosarcoma In most genetic studies no difference is made between peripheral, central chondrosarcoma and other subtypes. CGH revealed extensive genetic aberrations; gains of whole chromosomes or chromosome arms at 20q (32-38%), 20p (24-31%), and 14q23-qter (24-28%).

In a comparative study, 19 of 20 peripheral chondrosarcoma showed LOH at the loci for EXT, EXTL, 13q14, 17p14, 17p13, 9p21 and chromosome 10, while only 3 of 12 central chondrosarcoma did. In addition the ploidy status in peripheral chondrosarcoma showed wide variation (0,56-2,01), whereas central chondrosarcomas were predominantly periploid.

- **Dedifferentiated**
In the cartilaginous, as well as in the dedifferentiated part of the tumor an identical somatic 6 bp deletion in exon 7 of p53 and loss of the same copy on chromosome 13 had been found. These findings provide evidence for a common origin of both parts.
- **Mesenchymal**
A Robertsonian (13;21) translocation, der(13;21)(q10;q10), was detected in two cases, together with loss of material from chromosome 8 and 20 and gain of material from chromosome 8.
- **Clear cell**
No specific alterations have been reported.
- **Periosteal**
No specific alterations have been reported.

Cytogenetics **Osteogenic tumors**

Osteoid osteoma

Involvement of band 22q13 and loss of the distal part of arm 17q were detected in two out of three analyzed cases.

Osteoblastoma

No consistent aberrations have been detected in four cases, although clues are leading to deregulation of the cell cycle. No telomerase activity could be found.

Osteosarcoma

- **Conventional**
Although most osteosarcomas reveal complex chromosomal aberrations. no specific alterations have

been found. Amplification at 1q21-23 and 17p, together with co-amplification of 12q13-15 are frequently reported. In 14-27% of osteosarcomas, the [MDM2](#) gene is amplified and in 41% the [PRIM1](#) gene. Amplification of the [CDK4](#) gene is found in aggressive lesions. Chromosome 12p is differently amplified in low- versus high-grade osteosarcomas. The [MYC](#) gene has been shown to be amplified in 44% of osteosarcoma (8q24).

- Teleangiectatic
In contrast to the conventional osteosarcomas, no amplifications were detected in four cases. Complex chromosomal changes were found.
- Small cell
No genetic alterations have been reported on this subtype of osteosarcoma.
- Low grade central
The complex aberrations found in high-grade osteosarcomas could not be identified in these low-grade variants. Recurrent gains have been reported with minimal common regions at 12q13-14, 12p and 6p21.
- Secondary
In tumors developing in bone after radiation, chromosomal losses are more frequently observed than gains. Especially 1p, which is lost in 3% of the conventional type, is lost in 57% of the radiation-associated type.
In osteosarcomas secondary to Paget Disease of Bone (PDB) LOH is found at 18q, the locus for PDB.
- Parosteal
Ring chromosomes are reported as the sole alteration in parosteal osteosarcoma. Using FISH techniques, the [SAS](#), CDK4 and MDM2 genes are shown to be co-amplified with a minimal common region at 12q13-15 in the ring chromosomes.
- Periosteal
Complex chromosomal aberrations were reported.
- High grade surface
These lesions have not been studied as a specific subtype.

Cytogenetics **Fibrogenic tumors**

Desmoplastic fibroma of bone

Like in desmoid tumors, trisomies 8 and 20 are commonly found.

Fibrosarcoma of bone

No cytogenetic investigations on fibrosarcoma have been published.

Cytogenetics **Fibrohistiocytic tumors**

Histiocytoma of bone

Of the benign form of this tumor, no cytogenetic information is available. Its malignant counterpart shows LOH at chromosome 9p21-22, which has also been shown by CGH before.

Cytogenetics **Ewing sarcoma/Primitive neuroectodermal tumor**

The most common rearrangement (85%) in this tumor is translocation t(11;22)(q24;q12). Consequently, this leads to the fusion protein EWS/FLI1. Other translocations found in the Ewing Sarcoma gene are listed below:

fusion of	to	translocation	%
EWS	FLI1	t(11;22)(q24;q12)	85
EWS	ERG	t(21;22)(q22;q12)	10-15
EWS	ETV1	t(7;22)(p22;q12)	< 1
EWS	FEV	t(2;22)(q36;q12)	< 1
EWS	E1AF	t(17;22)(q21;q12)	< 1

Cytogenetics **Giant cell tumors**

Giant cell tumor

Reduction in length of the telomeres, as well as telomeric association, has been demonstrated in these tumors. Most commonly 11p, 13p, 14p, 15p, 19q, 20q and 21p are affected.

Cytogenetics **Notochordal tumors**

Chordoma

Nine of sixteen investigated chordomas were hypodiploid with a chromosome number ranging from 33 to 44. Chromosomes 3, 4, 10, and 13 are most commonly lost, and in half of the cases the following segments are lost up to the telomere: 1p31, 3p21, 3q21, 9p24, 17q11. Because LOH is found at band 1p36, a tumor suppressor gene is thought to exist on distal 1p.

Cytogenetics **Vascular tumors**

Haemangioma and related lesions

No cytogenetic investigations reported.

Angiosarcoma

An identical translocation of chromosomes 1 and 3 has been reported in two epithelioid haemangioendotheliomas.

Cytogenetics **Myogenic, lipogenic, neural and epithelial tumors**

Leiomyosarcoma of bone

5 Grade IIB tumors demonstrated a rate of genomic loss of 90%, whereas high micro satellite instability was not observed. Allelotyping revealed loss of pRb in the tumors. In addition, chromosomal loss was noticed in human telomerase subunit-linked markers.

Lipoma of bone

Only one case study is published about the benign form of this tumor. On its malignant counterpart, liposarcoma of bone, no genetic information has been published.

[Adamantinoma](#) and osteofibrous dysplasia

Cumulating evidence indicates that classic adamantinomas derive from their osteofibrous dysplasia (OFD)-like counterparts. OFD and adamantinoma show common cytogenetic abnormalities. In 15 cases of adamantinoma (n=11) and OFD (n=4) trisomies of chromosomes 7, 8, 12, 19 and 21 were detected. These findings further substantiate the clonal origin of OFD and the common histogenesis of OFD and adamantinoma.

Cytogenetics **Tumors of undefined neoplastic nature**

[Aneurysmal bone cyst](#)

Aneurysmal bone cysts can be primary or secondary to other bone lesions. Chromosome bands 16q22 and/or 17p13 ([USP6](#) gene) are non-randomly rearranged in ABC, regardless of tumor type (classic and solid) and of location (osseous and extraosseous). However, rearrangements are absent in secondary ABC. A recurrent t(16;17)(q22;p13) has been identified, but other chromosomal segments as translocation partner for each chromosome have been described.

Simple bone cyst

Only one case report describes structural rearrangements.

[Fibrous dysplasia](#)

Two fibrous dysplasia cases exhibited either a completely normal karyotype or single cell aberrations. Evidence that this lesion is neoplastic comes from the fact that clonal chromosomal aberrations have been found. In monostotic as well as polyostotic lesions activating [GNAS1](#)(20q13.2) mutations, known from the McCune-Albright syndrome, have been demonstrated.

Langerhans cell histiocytosis (LCH)

Studies of X-chromosome inactivation demonstrated that LCH is clonal.

Cytogenetics **Congenital and inherited syndromes**

[Beckwith-Wiedemann svndrome](#)

This syndrome, which is caused by heterogenic genetic changes on 11q15, is subject to genomic imprinting. 3 Beckwith-Wiedemann syndrome chromosome regions (BWSCR) have been identified: BWSC1 near INS/[IGF2](#), BWSC2 5 Mb proximal to BWSC1, and BWSC3 2 Mb even more proximal.

Enchondromatosis: Ollier disease and Maffucci syndrome
These syndromes are non-hereditary although a case of familial clustering has been reported. Mutations in the PTH receptor 1 have been found in two cases, but these results could not be confirmed in a larger series of 31 patients, suggesting that PTHR1 is not the culprit for enchondromatosis.

McCune-Albright syndrome
Mutations in the GNAS1 gene, located on 20q13, change the structure of the G-protein α -stimulatory subunit. Dysfunctions of the heterotrimeric G-protein complexes lead to this non-familial occurring disorder.

Multiple osteochondromatosis(MO)/[Hereditary Multiple Exostosis](#)
Mutations in one of the two exostosis (EXT) genes are responsible for this autosomal dominant syndrome. EXT1 is located at 8q24 and EXT2 at 11p11-p12. Most mutations are either nonsense, frame shift or splice-site mutations, leading to premature termination of the EXT proteins. This causes alteration of the gene products, which are functioning in the endoplasmic reticulum as transmembrane glycoproteins, and will affect the biosynthesis of heparan sulphate proteoglycans, leading to altered growth factor signaling.

Familial Paget Disease of Bone
Familial Paget Disease of Bone (PDB) demonstrates linkage to chromosome 18q. Some cases (PDB type 2) are caused by mutations in the [TNFRSF11A](#) gene on chromosome 18q22.1, which encodes RANK, a protein essential in osteoclast formation. The phenotype linked to chromosome 5q35 (PDB type 3) is caused by mutations in the SQSTM1 gene, the product of which is associated with the RANK pathway.

Bibliography

Telomeric associations and consistent growth factor overexpression detected in giant cell tumor of bone.

Schwartz HS, Butler MG, Jenkins RB, Miller DA, and Moses HL.
Cancer Genet Cytogenet 1991; 56: 263-276.
Medline [1756472](#)

Clonal karyotypic aberrations in enchondromas.

Bridge JA, Persons DL, Neff JR, and Bhatia P.
Cancer Detect Prev 1992; 16: 215-219.
Medline [1458512](#)

Biologic and clinical significance of cytogenetic and molecular cytogenetic abnormalities in benign and malignant cartilaginous lesions.

Bridge JA, Bhatia PS, Anderson JR, and Neff JR.
Cancer Genet Cytogenet 1993; 69: 79-90.
Medline [8402563](#)

Rearrangement of band q13 on both chromosomes 12 in a periosteal chondroma.

Mandahl N, Willen H, Rydholm A, Heim S, and Mitelman F.
Genes Chromosomes Cancer 1993; 6: 121-123
Medline [7680888](#)

Clonal chromosomal abnormalities in osteofibrous dysplasia. Implications for histopathogenesis and its relationship with adamantinoma.

Bridge JA, Dembinski A, De Boer J, Travis J, and Neff JR.
Cancer 1994; 73: 1746-1752.
Medline [8156503](#)

Langerhans'-cell histiocytosis (histiocytosis X). A clonal proliferative disease.

Willman C L, Busque L, Griffith BB, Favara BE, McClain KL, Duncan MH, and Gilliland DG.
N Engl J Med 1994; 331: 154-160.
Medline [8008029](#)

Mesenchymal chondrosarcoma - A cytogenetic, immunohistochemical and ultrastructural study.

Dobin SM, Donner LR, and Speights VO, Jr.
Cancer Genet Cytogenet 1995; 83: 56-60.
Medline [7656206](#)

DNA aberrations in the epithelial cell component of adamantinoma of long bones.

Hazelbag HM, Fleuren GJ, Cornelisse CJ, Van den Broek LJCM, Taminiau AHM, and Hogendoorn PCW.
Am J Pathol 1995; 147: 1770-1779.
Medline [7495301](#)

Gains and losses of DNA sequences in osteosarcomas by comparative genomic hybridization.

Tarkkanen M, Karhu R, Kallioniemi A, Elomaa I, Kivioja AH, Nevalainen J, Bohling T, Karaharju E, Hyytinen E, Knuutila S, and Kallioniemi O-P.
Cancer Res 1995; 55: 1334-1338.
Medline [7882332](#)

Positional cloning of a gene involved in hereditary multiple exostoses.

Wuyts W, Van Hul W, Wauters J, Nemtsova M, Reyniers E, Van Hul EV, De Boulle K, de Vries BB, Hendrickx J, Herrygers I, Bossuyt P, Balemans W, Fransen E, Vits L, Coucke P, Nowak NJ, Shows TB, Mallet L, van den Ouweland AM, McGaughan J,

Halley DJ, Willems PJ.
Hum Mol Genet 1996; 5: 1547-1557.
Medline [8894688](#)

Cytogenetic findings in 73 osteosarcoma specimens and a review of the literature.

Bridge JA, Nelson M, McComb E, McGuire MH, Rosenthal H, Vergara G, Maale GE, Spanier S, and Neff J R.
Cancer Genet Cytogenet 1997; 95: 74-87. Review.
Medline [9140456](#)

The pericentromeric inversion, inv (6)(p25q13), is a novel diagnostic marker in chondromyxoid fibroma.

Granter SR, Renshaw AA, Kozakewich HP, and Fletcher JA.
Mod Pathol 1998; 11: 1071-1074.
Medline [9831204](#)

Cytogenetic analysis of a scapular chondromyxoid fibroma.

Halbert AR, Harrison WR, Hicks MJ, Davino N, and Cooley LD.
Cancer Genet Cytogenet 1998; 104: 52-56.
Medline [9648559](#)

Evidence of an association between 6q13-21 chromosome aberrations and locally aggressive behavior in patients with cartilage tumors.

Sawyer JR, Swanson CM, Lukacs JL, Nicholas RW, North PE, and Thomas JR.
Cancer 1998; 82: 474-483.
Medline [9452264](#)

Significance of abnormalities of chromosomes 5 and 8 in chondroblastoma.

Swarts SJ, Neff JR, Johansson SL, Nelson M, and Bridge JA.
Clin Orthop 1998; 189-193.
Medline [9584382](#)

EXT-mutation analysis and loss of heterozygosity in sporadic and hereditary osteochondromas and secondary chondrosarcomas.

Bovee JV, Cleton-Jansen AM, Wuyts W, Caethoven G, Taminiau AH, Bakker E, Van Hul W, Cornelisse CJ, Hogendoorn PC.
Am J Hum Genet 1999; 65: 689-698.
Medline [10502322](#)

Loss of heterozygosity and DNA ploidy point to a diverging genetic mechanism in the origin of peripheral and central chondrosarcoma.

Bovee JV, Cleton-Jansen AM, Kuipers-Dijkshoorn NJ, van den Broek LJ, Taminiau AH, Cornelisse CJ, Hogendoorn PC.
Genes Chromosomes Cancer 1999; 26: 237-246.
Medline [10502322](#)

Molecular genetic characterization of both components of a dedifferentiated chondrosarcoma, with implications for its histogenesis.

Bovee JV, Cleton-Jansen AM, Rosenberg C, Taminiou AH, Cornelisse CJ, Hogendoorn PC.

J Pathol 1999; 189: 454-462.

Medline [10629543](#)

Recurrent chromosome aberrations in fibrous dysplasia of the bone: a report of the CHAMP study group. CHromosomes And MorPhology.

Dal Cin P, Sciot R, Brys P, De, Wever I, Dorfman H, Fletcher CD, Jonsson K, Mandahl N, Mertens F, Mitelman F, Rosai J, Rydholm A, Samson I, Tallini G, Van Den Berghe H, Vanni R, and Willen H.

Cancer Genet Cytogenet 2000; 122: 30-32.

Medline [11104029](#)

Clonal chromosome abnormalities in enchondromas and chondrosarcomas.

Gunawan B, Weber M, Bergmann F, Wildberger J, Niethard F U, and Fuzesi L.

Cancer Genet Cytogenet 2000; 120: 127-130.

Medline [10942802](#)

Molecular Cytogenetics in Ewing Tumors: Diagnostic and Prognostic Information.

Hattinger CM, Zoubek A, and Ambros PF.

Onkologie 2000; 23: 416-422.

Medline [11441235](#)

Malignant fibrous histiocytoma: inherited and sporadic forms have loss of heterozygosity at chromosome bands 9p21-22-evidence for a common genetic defect.

Martignetti, JA, Gelb, BD, Pierce H, Picci P, and Desnick R J.

Genes Chromosomes Cancer 2000; 27: 191-195.

Medline [10612808](#)

Recurrent anomalies of 6q25 in chondromyxoid fibroma.

Safar A, Nelson M, Neff JR, Maale GE, Bayani J, Squire J, and Bridge JA.

Hum Pathol 2000; 31: 306-311.

Medline [10746672](#)

Frequent loss of 9p21 (p16(INK4A)) and other genomic imbalances in human malignant fibrous histiocytoma.

Simons A., Schepens M., Jeuken J., Sprenger S, van de Zande G, Bjerkehagen B, Forus A, Weibolt V, Molenaar I, van den Berg E, Myklebost O, Bridge J, van Kessel, AG, and Suijkerbuijk R.

Cancer Genet Cytogenet 2000; 118: 89-98.

Medline [10748288](#)

EXT genes are differentially expressed in bone and cartilage during mouse embryogenesis.

Stickens D, Brown D, Evans GA.

Dev Dyn 2000; 218: 452-464.

Medline [9232192](#)

Analysis of the p16INK4, p14ARF, p15, TP53, and MDM2 genes and their prognostic implications in osteosarcoma and Ewing sarcoma.

Tsuchiya T, Sekine K, Hinohara S, Namiki T, Nobori T, and Kaneko Y

Cancer Genet Cytogenet 2000; 120: 91-98.

Medline [10942797](#)

Chromosome 9 alterations and trisomy 22 in central chondrosarcoma: a cytogenetic and DNA flow cytometric analysis of chondrosarcoma subtypes.

Bovée JVMG, Sciot R, Dal Cin P, Debiec-Rychter M, Van Zelderen-Bhola SL, Cornelisse CJ, and Hogendoorn PCW.

Diagn Mol Pathol 2001; 10: 228-236.

Medline [11763313](#)

Genome-wide analysis of sixteen chordomas by comparative genomic hybridization and cytogenetics of the first human chordoma cell line, U-CH1.

Scheil S, Bruderlein S, Liehr T, Starke H, Herms J, Schulte M, and Moller P.

Genes Chromosomes Cancer 2001; 32: 203-211.

Medline [11579460](#)

Cytogenetic and molecular cytogenetic evidence of recurrent 8q24.1 loss in osteochondroma.

Feely MG, Boehm AK, Bridge RS, Krallman PA, Neff JR, Nelson M, Bridge JA.

Cancer Genet Cytogenet 2002; 137: 102-107.

Medline [12393280](#)

WHO Classification of tumours. Pathology & Genetics. Tumours of Bone and Soft tissue.

Fletcher CDM, Unni KK, and Mertens F.

IARC Press: Lyon 2002.

Translocation der(13;21)(q10;q10) in Skeletal and Extraskelatal Mesenchymal Chondrosarcoma.

Naumann S, Krallman PA, Unni KK.

Mod Pathol 2002; 15: 572-576

Medline [12011263](#)

Recurring breakpoints of 1p13 approximately p22 in osteochondroma.

Sawyer JR, Thomas EL, Lukacs JL, Swanson CM, Ding Y, Parham DM, Thomas JR, Nicholas RW.

Cancer Genet Cytogenet 2002; 138: 102-106.

Medline [12505252](#)

Correlation between clinicopathological features and karyotype in 100 cartilaginous and chordoid tumours. A report from the Chromosomes and Morphology (CHAMP) Collaborative Study Group.

Tallini G, Dorfman H, Brys P, Dal Cin P, De Wever I, Fletcher CD, Jonson K, Mandahl N, Mertens F, Mitelman F, Rosai J, Rydholm A, Samson I, Sciot R, Van den Berghe H, Vanni R, Willen H.

J Pathol 2002; 196: 194-203.

Medline [11793371](#)

Genetic and epigenetic alterations in tumor progression in a dedifferentiated chondrosarcoma.

Ropke M, Boltze C, Neumann HW, Roessner A, Schneider-Stock R.

Pathol Res Pract 2003; 199: 437-444.

Medline [12924447](#)

Updates on the cytogenetics and molecular genetics of bone and soft tissue tumors: chondrosarcoma and other cartilaginous neoplasms.

Sandberg AA, Bridge JA.

Cancer Genetics and Cytogenetics 2003; 143: (1-31). Review.

Medline [12742153](#)

Multiple Osteochondromas: Clinicopathological and Genetic Spectrum and Suggestions for Clinical Management.

Hameetman L, Bovée JVMG, Taminiau AHM, Kroon HM, and Hogendoorn PCW.

Hereditary Cancer in Clinical Practice 2004; 2: 161-173. Review.

USP6 and CDH11 oncogenes identify the neoplastic cell in primary aneurysmal bone cysts and are absent in so-called secondary aneurysmal bone cysts.

Oliveira AM, Perez-Atayde AR, Inwards CY, Medeiros F, Derr V, Hsi BL, Gebhardt MC, Rosenberg AE, Fletcher JA.

Am J Pathol 2004; 165: 1773-1780.

Medline [15509545](#)

Cytogenetic distinction among benign fibro-osseous lesions of bone in children and adolescents: value of karyotypic findings in differential diagnosis.

Parham DM, Bridge JA, Lukacs JL, Ding Y, Tryka AF, Sawyer JR.

Pediatr Dev Pathol 2004; 7: 148-158.

Medline [1502206](#)

Enchondromatosis (Ollier disease, Maffucci syndrome) is not caused by the PTHR1 mutation p.R150C.

Rozeman LB, Sangiorgi L, Bruijn IH, Mainil-Varlet P, Bertoni F, Cleton-Jansen AM, Hogendoorn PC, and Bovée JVMG.

Hum Mutat 2004; 24: 466-473.

Medline [15523647](#)

Genetic instability in primary leiomyosarcoma of bone.

Verelst SJ, Hans J, Hanselmann RG, Wirbel RJ.

Hum Pathol 2004; 35: 1404-1412.

Medline [15668899](#)

REVIEW articles

automatic search in PubMed

Last year publications

automatic search in PubMed

Contributor(s)

Written 03-2005 Yvonne M Schrage, Judith VMG Bovée.

Citation

This paper should be referenced as such :

Schrage YM, Bovée JVMG . Bone tumors: an overview. Atlas Genet Cytogenet Oncol Haematol. March 2005 .

URL : <http://AtlasGeneticsOncology.org/Tumors/BoneTumorID5143.html>

© *Atlas of Genetics and Cytogenetics in Oncology and Haematology*
