

Bone: Fibrous dysplasia of the bone

Identity

Note In the major text-books on bone pathology, fibrous dysplasia (FD) is regarded as a non-neoplastic process. It is included in the chapter on "conditions that simulate a bone neoplasm" and it is defined elsewhere as a dysplastic disorder of bone.

Clinics and Pathology

Etiology Unknown

Clinics It is characterized by a broad clinical spectrum, varying from the more frequent solitary (monostatic) asymptomatic lesion to extensive and severe multifocal (polyostatic) lesions. The monostotic variant most commonly affects the ribs, femur and tibia of older children and young adults. The less frequent polyostotic type may be associated with endocrine abnormalities, skin hyperpigmentation and soft tissue myxoma (so-called Albright syndrome).

On radiographs, fibrous dysplasia presents as a well defined osteolytic lesion with benign features, centered within the medulla, and frequently bordered by a shell of reactive sclerosis. Depending on the relative proportion of osseous to fibrous tissue, the lesion can be completely lytic or, most typically, show a relatively high density, the so-called 'ground-glass' appearance in a least part of it. Especially in bones with small diameter and flat bones, fibrous dysplasia frequently expands the bone.

Pathology Microscopically, irregular trabeculae of woven bone are embedded in a moderately cellular fibrous matrix. The bony trabeculae have variable and irregular shapes. Osteoblastic rimming of the trabeculae is characteristically inconspicuous and some of the trabeculae seem to emerge from the surrounding fibrous background, suggesting a 'metaplastic' process. Benign lesions that histologically can be confused with FD are osteofibrous dysplasia of long bones and desmoplastic fibroma. Cases of FD with extensive cartilaginous differentiation may be mistaken histologically or radiologically for a benign chondroblastic tumor, although the adjacent fibro-osseous component allows this distinction. The most important differential diagnosis of FD is with low grade (well-differentiated) central [osteosarcoma](#). Rarely sarcomas can arise in FD and some of them occur in patients that had received radiation therapy (post-radiation sarcomas).

Treatment It should be conservative

Cytogenetics

Cytogenetics Clonal chromosome aberrations have been reported in 8 out of eleven Morphological cases of FD in which chromosomal analysis has been performed, suggesting that this entity is of a neoplastic nature. The only recurrent changes described so far in FD have been structural 12p13 aberrations (3 cases) and trisomy 2 (3 cases).

External links

[Orphanet](#) [Fibrous dysplasia of the bone](#)

Bibliography

Desmoplastic fibroma arising in fibrous dysplasia. Chromosomal analysis and review of the literature

Bridge JA, Rosenthal H, Sanger WG, Neff JR
Clin Orthop Rel Res 1989; 247:272-278
Medline [2676299](#)

Cytogenetic study of 249 consecutive patients examined for a bone tumor

Tarkkanen M, Kaipainen A, Karaharju E, Böhling T, Szymanska J, Helio H, Kivioja A, Elemaa I, Knuutila S
Cancer Genet Cytogenet 1993; 68:1-21
Medline [8330278](#)

Chromosome aberrations in fibrous dysplasia

Dal Cin P, Sciot R, Speleman F, Samson I, Laureys G, De Potter C, Meire F, Van Damme B, Van den Berghe H
Cancer Genet Cytogenet 1994; 77:114-117.
Medline [7954320](#)

Clonal structural chromosome aberrations in fibrous dysplasia

Mertens F, Albert A, Heim S, Lindholm J, Brosjö O, Mitelman, F, Mandahl N
Genes Chromosom Cancer 1994; 11:271-272
Medline [7533532](#)

Malignancies in fibrous dysplasia

Ruggeri P, Sim FH, Bond JR, Unni KK
Cancer 1994; 73:1411-1424

Dahlin's Bone Tumors. General Aspects and Data on 11.087 cases

Unni KK
5th ed. 1996 Lippincott-Raven Publishers, pp. 367-376.

Bone Tumors

Dorfman HD, Czerniak B
The Mosby Publisher 1998, pp. 441-481

Trisomies 8 and 20 characterize a subgroup of benign fibrous lesions arising in both soft tissue and bone

Bridge JA, Swartz SJ, Buresh C, Nelson M, Degenhardt Jm, Spanier S, Maale G, Meloni, A, Lynch JC, Neff JR
Am J Path 1999; 154:729-733
Medline [10079250](#)

Fibrous dysplasia and the short arm of chromosome 12

Dal Cin P, Bertoni F, Bacchini A, Hagemeyer A, Van Den Berghe H
Histopathology 1999; 34:279-280.
Medline [10217576](#)

Recurrent Chromosome Aberrations in Fibrous Dysplasia of the Bone: A Report of the CHAMP Study Group.

Dal Cin P, Sciot R, Brys P, De Wever I, Dorfman H, Fletcher CDM, Jonsson K, Mandahl N, Mertens F, Mitelman F, Rosai J, Rydholm A, Samson I, Tallini G, Van den Berghe H, Vanni R, Willen H

Cancer Genet Cytogenet 2000; 122:30-32.

Medline [11104029](#)

[REVIEW articles](#) *automatic search in PubMed*

[Last year publications](#) *automatic search in PubMed*

Contributor(s)

Written 06-2001 Paola Dal Cin

Citation

This paper should be referenced as such :

Dal Cin P . Bone: Fibrous dysplasia of the bone. Atlas Genet Cytogenet Oncol Haematol. June 2001 .

URL : <http://AtlasGeneticsOncology.org/Tumors/FibDysplasiaBoneID5093.html>

© *Atlas of Genetics and Cytogenetics in Oncology and Haematology*
